

AMVUTTRA® (vutrisiran) may help you

BRING LIFE'S MOMENTS **WITHIN REACH**

AMVUTTRA is the first and only injection given 4 times a year to treat the polyneuropathy of hereditary ATTR (hATTR) amyloidosis in adults

Patient portrayal.

In an 18-month study, patients receiving AMVUTTRA had significantly improved nerve function and quality of life compared to patients receiving placebo in a similar study.



Calms the symptoms
of polyneuropathy



Rapidly knocks
down TTR



Dosing once every 3 months
by a healthcare professional

Important Safety Information

What are the most important things I should know about AMVUTTRA?

AMVUTTRA can cause:

- **Low Vitamin A Levels**

Treatment with AMVUTTRA lowers the amount of vitamin A in your blood. Your doctor will tell you to take a vitamin A supplement every day. You should not take more than the amount of vitamin A recommended by your doctor. (Continued on page 5)

Please see additional **Important Safety Information** on page 16 and full **Prescribing Information**.

amvuttra
(vutrisiran) injection
25 mg/0.5 mL



hATTR amyloidosis is a rare and rapidly progressive condition

hATTR amyloidosis is caused by a variant, or change, in the transthyretin (TTR) gene



It runs in the family

- If one parent has hATTR amyloidosis, **each child will have a 50% chance** of inheriting the genetic variant that causes the condition
- A family member may inherit the TTR gene variant, but having the variant does not necessarily mean they will develop hATTR amyloidosis
- The age that symptoms typically appear ranges from **the mid-20s to the mid-60s**



Difficult to recognize

- hATTR amyloidosis is often misdiagnosed because its symptoms resemble those of other conditions
- If your doctor suspects that you have hATTR amyloidosis, they may order further tests to make a diagnosis. Some common tests include^a:
 - Genetic testing
 - Tissue biopsy


^aThis is not a complete list of tests that can help diagnose hATTR amyloidosis.



Being diagnosed with hATTR amyloidosis

- It can be difficult to come to terms with a diagnosis of hATTR amyloidosis, and you may have questions about the disease
- It's important to know that you're not alone. There are resources available to help you understand your diagnosis

How hATTR amyloidosis develops



Talk to your doctor if you think you are experiencing symptoms of hATTR amyloidosis.

Symptoms of hATTR amyloidosis

hATTR amyloidosis can cause symptoms that affect several parts of the body, such as the nerves, heart, and digestive system, which can worsen quickly if left untreated.

Nerve damage that affects sensation, movement, strength, the digestive system, and other bodily functions may be referred to **as polyneuropathy, and can cause the following:**

Symptoms related to nerves in hands and feet, arms and legs:

- Burning pain
- Carpal tunnel syndrome
- Decreased reflexes
- Difficulty walking
- Temperature sensitivity loss
- Tingling and/or numbness
- Weakness



Symptoms related to the digestive system:

- Diarrhea
- Feeling full quickly
- Nausea
- Severe constipation
- Unintentional weight loss
- Vomiting



Additional symptoms:

- Abnormal sweating
- Difficulty urinating
- Dizziness upon standing
- Recurrent urinary tract infections
- Sexual dysfunction



Other symptoms of hATTR amyloidosis include:


- Blurred or spotty vision
- Floaters in eyes
- Glaucoma
- Kidney dysfunction
- Heart failure and other cardiac symptoms such as:
 - Abnormal heart rhythms (atrial fibrillation)
 - Fatigue
 - Leg swelling (edema)
 - Palpitations
 - Shortness of breath

This is not a complete list of symptoms that may be experienced with hATTR amyloidosis.

AMVUTTRA® (vutrisiran) does not treat all of the symptoms of hATTR amyloidosis.

AMVUTTRA[®] rapidly knocks down TTR

AMVUTTRA works with your body's natural system to rapidly knock down TTR. It decreases the amount of TTR protein made in the liver, which leads to fewer harmful amyloid deposits being formed.



AMVUTTRA starts reducing TTR protein after the first dose, which lasts with continued treatment. TTR protein was reduced by an average of 88% over 18 months in a clinical study of adults treated with AMVUTTRA.

Important Safety Information

What are the most important things I should know about AMVUTTRA?

AMVUTTRA can cause:

- **Low Vitamin A Levels (cont.)**

Low vitamin A levels can affect vision. If you have problems with your vision (e.g., night blindness) while taking AMVUTTRA, talk to your doctor. Your doctor may refer you to an eye specialist.

Please see additional [Important Safety Information](#) on page 16 and full [Prescribing Information](#).



**“With this disease, you definitely need a holistic and team approach
so you can see the complete picture.”**

—AL, treated with AMVUTTRA® (vutrisiran), and **DEBORAH**, his spouse and caregiver

Please see [Important Safety Information](#) on page 16 and full [Prescribing Information](#).

How AMVUTTRA[®] was studied

The safety and efficacy of AMVUTTRA were evaluated in an 18-month study of 164 adult patients with polyneuropathy caused by hATTR amyloidosis.

The clinical study of AMVUTTRA focused on changes in **nerve function** and **quality of life**.

The effects of AMVUTTRA on other aspects of the disease, including **walking speed**, **nutritional health**, and **ability to perform activities of daily living**, were also evaluated.



The efficacy of AMVUTTRA was evaluated using multiple assessments



These assessments were performed at 9 months and again at 18 months



Efficacy was determined by comparing results from the following groups of patients with polyneuropathy caused by hATTR amyloidosis:

- Patients treated with AMVUTTRA
- Patients from a similar study who were given a placebo^a

^aPlacebo: treatment without any active medication.

Important Safety Information

What are the common side effects of AMVUTTRA?

The most common side effects of AMVUTTRA are pain in the arms or legs, pain in the joints (arthralgia), shortness of breath (dyspnea), and low vitamin A levels.


These are not all the possible side effects of AMVUTTRA. Talk to your doctor about side effects that you experience. You are encouraged to report negative side effects of prescription drugs to the U.S. Food and Drug Administration (FDA). Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see additional **Important Safety Information** on page 16 and full **Prescribing Information**.


AMVUTTRA[®] works to calm the symptoms of polyneuropathy

Patients treated with AMVUTTRA showed significant improvement in nerve function and quality of life at 9 months, and continued to improve throughout the 18-month clinical study, compared with those who received placebo in a similar study.

AMVUTTRA significantly improved
nerve function and quality of life



Patients who received placebo
got worse



Nerve function was assessed using a scale called mNIS+7 that measured strength and sensation in the hands, feet, arms, and legs; reflexes; and blood pressure upon standing.



Quality of life was evaluated by the Norfolk QoL-DN questionnaire that asked patients about the severity of their polyneuropathy symptoms, how often they experienced them, and what impact they felt they had on their daily lives.

Please see [Important Safety Information](#) on page 16 and full [Prescribing Information](#).



“After a discussion with my doctor, I thought AMVUTTRA would be the best treatment for me.”

—STEVE, treated with AMVUTTRA® (vutrisiran)

Important Safety Information

What are the most important things I should know about AMVUTTRA?

AMVUTTRA can cause:

- **Low Vitamin A Levels**

Treatment with AMVUTTRA lowers the amount of vitamin A in your blood. Your doctor will tell you to take a vitamin A supplement every day. You should not take more than the amount of vitamin A recommended by your doctor. (Continued on page 11)

Please see additional **Important Safety Information** on page 16 and full **Prescribing Information**.



“My fight to get the correct diagnosis was a way to show my loved ones and the community the importance of being an advocate for yourself and for your own health.”

—JOHN, treated with AMVUTTRA® (vutrisiran)

Please see [Important Safety Information](#) on page 16 and full [Prescribing Information](#).

Patients continued to improve throughout treatment

Results from the study^a were evaluated to determine how many patients regained some nerve function or reported better quality of life from the start of treatment.

AT 18 MONTHS:

48% of patients treated with AMVUTTRA[®] **regained some nerve function** from the start of treatment, compared with **4% of those who received placebo**

- For the patients who received AMVUTTRA and did not regain some nerve function, progression of their neuropathy was slowed compared with those who received placebo

57% of patients treated with AMVUTTRA **reported better quality of life** from the start of treatment, compared with **10% of those who received placebo**

^aTotal number of patients in the 18-month study: 164.

Ask your doctor if AMVUTTRA is right for you.

Important Safety Information

What are the most important things I should know about AMVUTTRA?

AMVUTTRA can cause:

- **Low Vitamin A Levels (cont.)**

Low vitamin A levels can affect vision. If you have problems with your vision (e.g., night blindness) while taking AMVUTTRA, talk to your doctor. Your doctor may refer you to an eye specialist.

Please see additional [Important Safety Information](#) on page 16 and full [Prescribing Information](#).

Other improvements patients experienced

The effects of AMVUTTRA[®] on other aspects of the disease, including **walking speed, nutritional health, and ability to perform activities of daily living**, were also evaluated at 18 months and compared with the effects of placebo in a similar study.

Compared with patients who received placebo, patients treated with AMVUTTRA:



MAINTAINED A BETTER WALKING SPEED

- Evaluated using the 10-meter walk test, a stopwatch-timed measure of a patient's walking speed over 10 meters



IMPROVED IN NUTRITIONAL HEALTH AFTER 1 DOSE

- Evaluated using modified body mass index, an assessment of height, weight, and the balance of fluids in the body, and was maintained with continued dosing at 18 months



WERE BETTER ABLE TO PERFORM COMMON DAILY ACTIVITIES

- Evaluated using a questionnaire called R-ODS to assess ability to perform activities
- Common daily activities included:



Showering



Brushing
teeth



Dressing
upper body



Going
shopping



Doing the
dishes



Walking up a
flight of stairs

Please see [Important Safety Information](#) on page 16 and full [Prescribing Information](#).



**“I’d like other patients battling this disease to know
that you don’t have to let it define you.”**

—KELLY, treated with AMVUTTRA® (vutrisiran)

Important Safety Information

What are the common side effects of AMVUTTRA?

The most common side effects of AMVUTTRA are pain in the arms or legs, pain in the joints (arthralgia), shortness of breath (dyspnea), and low vitamin A levels.

These are not all the possible side effects of AMVUTTRA. Talk to your doctor about side effects that you experience. You are encouraged to report negative side effects of prescription drugs to the U.S. Food and Drug Administration (FDA). Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see additional **Important Safety Information** on page 16 and full **Prescribing Information**.

Possible side effects

- The most common side effects in patients treated with AMVUTTRA[®] during the clinical study were pain in the arms or legs (15%), pain in the joints (11%), shortness of breath (7%), and low vitamin A levels (7%)
- Injection site reactions were mild in severity, decreased over time, and occurred in 5 patients (4%)

Talk to your doctor if you have any questions
about treatment with AMVUTTRA.

Important Safety Information

What are the most important things I should know about AMVUTTRA?

AMVUTTRA can cause:

- **Low Vitamin A Levels**

Treatment with AMVUTTRA lowers the amount of vitamin A in your blood. Your doctor will tell you to take a vitamin A supplement every day. You should not take more than the amount of vitamin A recommended by your doctor. (Continued on back cover)

Please see additional [Important Safety Information](#) on page 16 and full [Prescribing Information](#).

Dosing only 4 times a year by a healthcare professional




Receiving AMVUTTRA[®]

- AMVUTTRA is given as a subcutaneous injection once every 3 months
- Subcutaneous injections are given under the skin, and the process may take a few minutes

Treatment site options

- You can work with your doctor to choose where you receive AMVUTTRA—your doctor's office, a local clinic, or your own home (depending on insurance)

AMVUTTRA can be given in the following areas of the body:



With AMVUTTRA treatment, no premedications or laboratory monitoring are required.

Please see [Important Safety Information](#) on page 16 and full [Prescribing Information](#).

Indication and Important Safety Information



Indication

What is AMVUTTRA® (vutrisiran)?

AMVUTTRA is a prescription medicine that treats the polyneuropathy caused by an illness called hereditary transthyretin-mediated amyloidosis (hATTR amyloidosis). AMVUTTRA is used in adults only.

Important Safety Information

What are the most important things I should know about AMVUTTRA?

AMVUTTRA can cause:

- **Low Vitamin A Levels**

Treatment with AMVUTTRA lowers the amount of vitamin A in your blood. Your doctor will tell you to take a vitamin A supplement every day. You should not take more than the amount of vitamin A recommended by your doctor.

Low vitamin A levels can affect vision. If you have problems with your vision (e.g., night blindness) while taking AMVUTTRA, talk to your doctor. Your doctor may refer you to an eye specialist.

What are the common side effects of AMVUTTRA?

The most common side effects of AMVUTTRA are pain in the arms or legs, pain in the joints (arthralgia), shortness of breath (dyspnea), and low vitamin A levels.

These are not all the possible side effects of AMVUTTRA. Talk to your doctor about side effects that you experience. You are encouraged to report negative side effects of prescription drugs to the U.S. Food and Drug Administration (FDA). Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

For additional information about AMVUTTRA, please see full [Prescribing Information](#).

Support and resources

Connect with an Alnylam Educator



An Alnylam Patient Education Liaison (PEL) can provide education to help you and your family better understand the disease and answer your questions about AMVUTTRA[®].

Alnylam PELs do not offer medical advice. Please talk to your doctor for specific concerns.

Visit www.amvuttraPEL.com to connect with an Alnylam Educator

Get patient support services through Alnylam Assist[®]



Alnylam Assist is a patient support program designed to help get you started on treatment, understand your insurance coverage, determine eligibility for financial assistance, and provide you and your family with ongoing support services in your treatment journey with AMVUTTRA.

An Alnylam Case Manager will work with you to begin treatment with and maintain access to AMVUTTRA.

 **1-833-256-2748**
Monday–Friday, 8AM–6PM

Financial assistance programs^a may include:

Alnylam Assist Commercial Copay Program^b covers certain out-of-pocket costs for eligible patients with commercial insurance.

Alnylam Assist Patient Assistance Program (PAP) provides AMVUTTRA at no cost to eligible patients, primarily the uninsured, who meet specified financial criteria.



To learn more, visit
www.AlnylamAssist.com.

^aPatients must meet specified eligibility criteria to qualify for assistance. Alnylam reserves the right to make eligibility determinations and to modify or discontinue any program at any time.

^bPatients with Medicare, Medicaid, or other government-sponsored insurance are not eligible for the Alnylam Assist Commercial Copay Program. Out-of-pocket costs for the administration of AMVUTTRA will not be covered for patients residing where it is prohibited by law or where otherwise restricted.

AMVUTTRA® (VUTRISIRAN) IS THE FIRST AND ONLY INJECTION GIVEN 4 TIMES A YEAR to treat the polyneuropathy of hATTR amyloidosis in adults



In an 18-month study, patients receiving AMVUTTRA had significantly improved nerve function and quality of life compared to patients receiving placebo in a similar study.



**Calms the symptoms
of polyneuropathy**



**Rapidly knocks
down TTR**



**Dosing once every 3 months
by a healthcare professional**

More than 1000 adult patients in the US have chosen AMVUTTRA to treat the polyneuropathy caused by hATTR amyloidosis.*

Learn more at
www.amvuttra.com

*As of October 2023.

Important Safety Information

What are the most important things I should know about AMVUTTRA?

AMVUTTRA can cause:

- **Low Vitamin A Levels (cont.)**

Low vitamin A levels can affect vision. If you have problems with your vision (e.g., night blindness) while taking AMVUTTRA, talk to your doctor. Your doctor may refer you to an eye specialist.

Please see additional [Important Safety Information](#) on page 16 and full [Prescribing Information](#).



AMVUTTRA, Alnylam Assist, and their associated logos are trademarks of Alnylam Pharmaceuticals, Inc.
© 2024 Alnylam Pharmaceuticals, Inc. All rights reserved.
AMV-USA-00298-V3



HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use AMVUTTRA™ safely and effectively. See full prescribing information for AMVUTTRA.

AMVUTTRA (vutrisiran) injection, for subcutaneous use
Initial U.S. Approval: 2022

INDICATIONS AND USAGE

AMVUTTRA is a transthyretin-directed small interfering RNA indicated for the treatment of the polyneuropathy of hereditary transthyretin-mediated amyloidosis in adults. (1)

DOSAGE AND ADMINISTRATION

- The recommended dosage of AMVUTTRA is 25 mg administered by subcutaneous injection once every 3 months. (2.1)
- AMVUTTRA is for subcutaneous use only and should be administered by a healthcare professional. (2.2)

DOSAGE FORMS AND STRENGTHS

Injection: 25 mg/0.5 mL in a single-dose prefilled syringe. (3)

CONTRAINDICATIONS

None. (4)

WARNINGS AND PRECAUTIONS

Reduced serum vitamin A levels and recommended supplementation: Supplement with the recommended daily allowance of vitamin A. Refer to an ophthalmologist if ocular symptoms suggestive of vitamin A deficiency occur. (5.1)

ADVERSE REACTIONS

The most common adverse reactions (≥5%) were pain in extremity, arthralgia, dyspnea, and vitamin A decreased. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Alnylam Pharmaceuticals at 1-877-256-9526 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 2/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

2.2 Administration Instructions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Reduced Serum Vitamin A Levels and Recommended Supplementation

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Immunogenicity

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

8.7 Hepatic Impairment

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

AMVUTTRA is indicated for the treatment of the polyneuropathy of hereditary transthyretin-mediated amyloidosis in adults.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

The recommended dosage of AMVUTTRA is 25 mg administered by subcutaneous injection once every 3 months [see [Dosage and Administration \(2.2\)](#)].


Missed Dose

If a dose is missed, administer AMVUTTRA as soon as possible. Resume dosing every 3 months from the most recently administered dose.

2.2 Administration Instructions

AMVUTTRA is for subcutaneous use only and should be administered by a healthcare professional.

Syringe Appearance Before and After Use



Preparation and Administration

1. Prepare the syringe

If stored cold, allow the syringe to warm to room temperature for 30 minutes prior to use.

Remove the syringe from the packaging by gripping the syringe body.

Do not touch the plunger rod until ready to inject.

Visually inspect the drug solution for discoloration and particulate matter prior to administration. AMVUTTRA is a sterile, preservative-free, clear, colorless-to-yellow solution. **Do not** use if it contains particulate matter or if it is cloudy or discolored.

Check the following:

- Syringe is not damaged, such as cracked or leaking
- Needle cap is attached to the syringe
- Expiration date on syringe label

Do not use the syringe if any issues are found while checking the syringe.


2. *Choose and prepare the injection site*

Choose an injection site from the following areas: the abdomen, thighs, or upper arms.

Avoid the following:

- 5-cm area around the navel
- Scar tissue or areas that are reddened, inflamed, or swollen

Clean the chosen injection site.



3. *Prepare the syringe for injection*

Hold the syringe body with one hand. Pull the needle cap straight off with other hand and dispose of needle cap immediately. It is normal to see a drop of liquid at the tip of the needle.

Do not touch the needle or let it touch any surface.

Do not recap the syringe.


Do not use the syringe if it is dropped.



4. *Perform the injection*


Pinch the cleaned skin.

Fully insert the needle into the pinched skin at a 45°-90° angle.




Inject all of the medication.

Push the plunger rod as far as it will go to administer the dose and activate the needle shield.



Release the plunger rod to allow the needle shield to cover the needle.

Do not block plunger rod movement.



5. *Dispose of the syringe*

Immediately dispose of the used syringe into a sharps container.

3 DOSAGE FORMS AND STRENGTHS

Injection: 25 mg/0.5 mL of vutrisiran as a clear, colorless-to-yellow solution in a single-dose prefilled syringe.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Reduced Serum Vitamin A Levels and Recommended Supplementation

AMVUTTRA treatment leads to a decrease in serum vitamin A levels [see [Adverse Reactions \(6.1\)](#) and [Clinical Pharmacology \(12.2\)](#)].

Supplementation at the recommended daily allowance of vitamin A is advised for patients taking AMVUTTRA. Higher doses than the recommended daily allowance of vitamin A should not be given to try to achieve normal serum vitamin A levels during treatment with AMVUTTRA, as serum vitamin A levels do not reflect the total vitamin A in the body.

Patients should be referred to an ophthalmologist if they develop ocular symptoms suggestive of vitamin A deficiency (e.g., night blindness).

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are discussed in greater detail in other sections of the labeling:

- Reduced Serum Vitamin A Levels and Recommended Supplementation [see [Warnings and Precautions \(5.1\)](#)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of AMVUTTRA cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In Study 1 [see [Clinical Studies \(14\)](#)], a total of 122 patients with polyneuropathy caused by hereditary transthyretin-mediated amyloidosis (hATTR amyloidosis) received AMVUTTRA. Of these, 118 patients received at least 18 months of treatment. The mean duration of treatment was 18.8 months (range: 1.7 to 19.4 months). The median patient age at baseline was 60 years and 65% of the patients were male. Seventy percent of AMVUTTRA-treated patients were Caucasian, 17% were Asian, 3% were Black, and 9% were reported as Other. Forty-four percent of patients had the Val30Met mutation in the transthyretin gene; the remaining patients had one of 21 other mutations. At baseline, 70% of patients were in Stage 1 of the disease and 30% were in Stage 2.

The most common adverse reactions (at least 5%) were pain in extremity, arthralgia, dyspnea, and vitamin A decreased (see [Table 1](#)).

In Study 1, patients were instructed to take the recommended daily allowance of vitamin A [see [Warnings and Precautions \(5.1\)](#)]. Seventy-four percent of patients treated with AMVUTTRA had normal vitamin A levels at baseline, and 98% of those with a normal baseline developed low vitamin A levels. In some cases, the decreased vitamin A level was reported as an adverse reaction (see [Table 1](#)).

Table 1: Adverse Reactions Reported in at least 5% of Patients Treated with AMVUTTRA (Study 1)

Adverse Reaction	AMVUTTRA
	N=122 %
Pain in extremity*	15
Arthralgia*	11
Dyspnea*	7
Vitamin A decreased†	7
*Comprised of several similar terms	
†Percentage only reflects those reported as an adverse reaction	

Two serious adverse reactions of atrioventricular (AV) heart block (1.6%) occurred in patients treated with AMVUTTRA, including one case of complete AV block.

Injection site reactions were reported in 5 (4%) patients treated with AMVUTTRA. Reported symptoms included bruising, erythema, pain, pruritus, and warmth. Injection site reactions were mild and transient.

6.2 Immunogenicity

As with all oligonucleotides, there is potential for immunogenicity. The detection of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies in the studies described below with the incidence of antibodies in other studies or to other products may be misleading.

In Study 1, 3 (2.5%) patients treated with AMVUTTRA developed anti-drug antibodies. Although anti-drug antibody development was not found to affect the pharmacokinetics, safety, or efficacy of AMVUTTRA in these patients, the available data are too limited to make definitive conclusions.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no available data on AMVUTTRA use in pregnant women to inform a drug-associated risk of adverse developmental outcomes. AMVUTTRA treatment leads to a decrease in serum vitamin A levels, and vitamin A supplementation is advised for patients taking AMVUTTRA. Vitamin A is essential for normal embryofetal development; however, excessive levels of vitamin A are associated with adverse developmental effects. The effects on the fetus of a reduction in maternal serum TTR caused by AMVUTTRA and of vitamin A supplementation are unknown [see [Warnings and Precautions \(5.1\)](#) and [Clinical Pharmacology \(12.2\)](#)].

In animal studies, subcutaneous administration of vutrisiran to pregnant rats resulted in developmental toxicity (reduced fetal body weight and embryofetal mortality) at doses associated with maternal toxicity (see [Data](#)).

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. The background risk of major birth defects and miscarriage for the indicated population is unknown.

Data

Animal Data

Subcutaneous administration of vutrisiran (0, 3, 10, or 30 mg/kg/day) to pregnant rats during the period of organogenesis resulted in embryofetal mortality at the high dose and reduced fetal body weight at the mid and high doses, which were associated with maternal toxicity.

Subcutaneous administration of vutrisiran (0, 3, 10, or 30 mg/kg/day) to pregnant rabbits resulted in no adverse effects on embryofetal development.

Subcutaneous administration of vutrisiran (0, 5, 10, or 20 mg/kg) to pregnant rats every 6 days throughout pregnancy and lactation resulted in no adverse developmental effects on the offspring.

8.2 Lactation

Risk Summary

There is no information regarding the presence of vutrisiran in human milk, the effects on the breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for AMVUTTRA and any potential adverse effects on the breastfed infant from AMVUTTRA or from the underlying maternal condition.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

8.5 Geriatric Use

No dose adjustment is required in patients ≥ 65 years of age [see [Clinical Pharmacology \(12.3\)](#)]. A total of 46 (38%) patients ≥ 65 years of age, including 7 (6%) patients ≥ 75 years of age, received AMVUTTRA in Study 1. No overall differences in safety or effectiveness were observed between these patients and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

8.6 Renal Impairment

No dose adjustment is recommended in patients with mild or moderate renal impairment (estimated glomerular filtration rate [eGFR] ≥ 30 to < 90 mL/min/1.73 m²) [see [Clinical Pharmacology \(12.3\)](#)]. AMVUTTRA has not been studied in patients with severe renal impairment or end-stage renal disease.



8.7 Hepatic Impairment

No dose adjustment is recommended in patients with mild hepatic impairment (total bilirubin $\leq 1 \times$ ULN and AST $>1 \times$ ULN, or total bilirubin >1.0 to $1.5 \times$ ULN and any AST) [see [Clinical Pharmacology \(12.3\)](#)]. AMVUTTRA has not been studied in patients with moderate or severe hepatic impairment.

11 DESCRIPTION

AMVUTTRA contains vutrisiran, a chemically modified double-stranded small interfering ribonucleic acid (siRNA) that targets mutant and wild-type transthyretin (TTR) messenger RNA (mRNA) and is covalently linked to a ligand containing three *N*-acetylgalactosamine (GalNAc) residues to enable delivery of the siRNA to hepatocytes.

The structural formula of vutrisiran sodium is presented below.



The molecular formula of vutrisiran sodium is $C_{530}H_{672}F_9N_{171}Na_{43}O_{323}P_{43}S_6$ with a molecular weight of 17,290 Da. The molecular formula of the free acid is $C_{530}H_{715}F_9N_{171}O_{323}P_{43}S_6$ with a molecular weight of 16,345 Da.

AMVUTTRA is supplied as a sterile, preservative-free, clear, colorless-to-yellow solution for subcutaneous injection. Each 0.5 mL of solution contains 25 mg of vutrisiran (equivalent to 26.5 mg vutrisiran sodium), 0.2 mg sodium phosphate monobasic dihydrate, 0.7 mg sodium phosphate dibasic dihydrate, 3.2 mg sodium chloride, water for injection, and sodium hydroxide and/or phosphoric acid to adjust the pH to ~ 7 .

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Vutrisiran is a double-stranded siRNA-GalNAc conjugate that causes degradation of mutant and wild-type TTR mRNA through RNA interference, which results in a reduction of serum TTR protein and TTR protein deposits in tissues.

12.2 Pharmacodynamics

In Study 1 [see [Clinical Studies \(14\)](#)], following administration of the recommended AMVUTTRA dosage every 3 months to patients with hATTR amyloidosis, vutrisiran reduced mean serum TTR at steady state by 83%. Similar TTR reductions were observed regardless of Val30Met genotype status, weight, sex, age, or race.

Vutrisiran also reduced the mean steady state serum vitamin A by 62% over 9 months [see [Warnings and Precautions \(5.1\)](#)].

Cardiac Electrophysiology

At a dose 12 times the recommended dosage of 25 mg once every three months, AMVUTTRA does not prolong the QT interval to any clinically relevant extent.

12.3 Pharmacokinetics

The pharmacokinetic (PK) properties of AMVUTTRA were evaluated following a single dose in healthy subjects and multiple doses in patients with hATTR amyloidosis, as summarized in [Table 2](#).

Table 2: Pharmacokinetic Parameters of Vutrisiran

	Vutrisiran
General Information	
Dose Proportionality	Vutrisiran C _{max} showed dose proportional increase while AUC _{last} and AUC _{inf} were slightly more than dose proportional following single subcutaneous doses ranging from 5 to 300 mg (i.e., 0.2 to 12 times the recommended dose)
Accumulation	No accumulation of vutrisiran was observed in plasma after repeated every 3 months dosage*
Absorption	
T_{max} [Median (Range)]	4 (0.17, 12.0) hours [†]
Distribution	
Estimated Vd/F (%RSE)	10.1 (5.8) L [‡]
Protein Binding	80% [§]
Organ Distribution	Vutrisiran distributes primarily to the liver after subcutaneous dosing
Elimination	
Half-Life [Median (Range)]	5.2 (2.2, 6.4) hours [†]
Apparent Clearance [Median (Range)]	21.4 (19.8, 30) L/hour [†]
Metabolism	
Primary Pathway	Vutrisiran is metabolized by endo- and exonucleases to short nucleotide fragments of varying sizes within the liver
Excretion	
Primary Pathway	The mean fraction of unchanged vutrisiran eliminated in urine was approximately 19.4% at the recommended dose of 25 mg. The mean renal clearance of vutrisiran ranged from 4.5 to 5.7 L/hour [¶]
AUC_{inf} = area under the concentration-time curve from the time of dosing extrapolated to infinity; AUC_{last} = area under the concentration-time curve from the time of dosing to the last measurable concentration; C_{max} = maximum plasma concentration; CV = coefficient of variation; RSE = relative standard error; T_{max} = time to maximum concentration; Vd/F = apparent volume of distribution *After 25 mg every 3 months dosage in hATTR amyloidosis patients [†]After 25 mg single dose in healthy subjects [‡]Based on population PK model estimation [§]Vutrisiran plasma protein binding was concentration-dependent and decreased with increasing vutrisiran concentrations (from 78% at 0.5 mcg/mL to 19% at 50 mcg/mL) [¶]After single subcutaneous vutrisiran dose from 5 to 300 mg (i.e., 0.2 to 12 times the recommended dose) in healthy subjects	

Specific Populations

No clinically significant differences in the pharmacokinetics of vutrisiran were observed based on age, sex, race, mild and moderate renal impairment (eGFR ≥ 30 to < 90 mL/min/1.73 m²), or mild hepatic impairment (total bilirubin ≤ 1 x ULN and AST > 1 x ULN, or total bilirubin > 1.0 to 1.5 x ULN and any AST). Vutrisiran has not been studied in patients with severe renal impairment, end-stage renal disease, moderate or severe hepatic impairment, or in patients with prior liver transplant.

Drug Interaction Studies

No clinical drug-drug interaction studies have been performed with vutrisiran. In vitro studies suggest that vutrisiran is not a substrate or inhibitor of cytochrome P450 enzymes. Vutrisiran is not expected to cause drug-drug interactions by inducing CYP enzymes or modulating the activities of drug transporters.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Subcutaneous administration of vutrisiran to male rats (0, 4, 7.5, or 15 mg/kg once every 4 weeks or 15 mg/kg once every 12 weeks) for 99 weeks and to female rats (0, 6, 12.5, or 25 mg/kg once every 4 weeks or 25 mg/kg once every 12 weeks) for 86-87 weeks resulted in no increase in tumors.

Mutagenesis

Vutrisiran was negative for mutagenicity in in vitro (bacterial mutagenicity, chromosomal aberration in human blood peripheral lymphocytes) and in vivo (rat bone marrow micronucleus) assays.

Impairment of Fertility

Subcutaneous administration of vutrisiran (0, 15, 30, or 70 mg/kg/week) to male and female rats prior to and during mating and continuing in females to gestation day 6 resulted in no adverse effects on fertility or reproductive performance.

14 CLINICAL STUDIES

The efficacy of AMVUTTRA was evaluated in a randomized, open-label clinical trial in adult patients with polyneuropathy caused by hATTR amyloidosis (Study 1; NCT03759379). Patients were randomized 3:1 to receive 25 mg of AMVUTTRA subcutaneously once every 3 months (N=122), or 0.3 mg/kg patisiran intravenously every 3 weeks (N=42) as a reference group. Ninety-seven percent of AMVUTTRA-treated patients and 93% of patisiran-treated patients completed at least 9 months of the assigned treatment.

Efficacy assessments were based on a comparison of the AMVUTTRA arm of Study 1 with an external placebo group in another study (NCT01960348) composed of a comparable population of adult patients with polyneuropathy caused by hATTR amyloidosis.

The primary efficacy endpoint was the change from baseline to Month 9 in modified Neuropathy Impairment Score +7 (mNIS+7). The mNIS+7 is an objective assessment of neuropathy and comprises the NIS and Modified +7 composite scores. In the version of the mNIS+7 used in the trial, the NIS objectively measures deficits in cranial nerve function, muscle strength, and reflexes, and the +7 assesses postural blood pressure, quantitative sensory testing, and peripheral nerve electrophysiology. The mNIS+7 has a total score range from 0 to 304 points, with higher scores representing a greater severity of disease.

The clinical meaningfulness of effects on the mNIS+7 was assessed by the change from baseline to Month 9 in Norfolk Quality of Life-Diabetic Neuropathy (QoL-DN) total score. The Norfolk QoL-DN scale is a patient-reported assessment that evaluates the subjective experience of neuropathy in the following domains: physical functioning/large fiber neuropathy, activities of daily living, symptoms, small fiber neuropathy, and autonomic neuropathy. The Norfolk QoL-DN has a total score range from -4 to 136, with higher scores representing greater impairment.

Additional endpoints were gait speed, as measured by the 10-meter walk test (10MWT), and modified body mass index (mBMI).

Treatment with AMVUTTRA in Study 1 resulted in statistically significant improvements in the mNIS+7, Norfolk QoL-DN total score, and 10-meter walk test at Month 9 compared to placebo in the external study ($p < 0.001$) [Table 3, Figure 1, and Figure 3]. The distributions of changes in mNIS+7 and Norfolk QoL-DN total scores from baseline to Month 9 by percent of patients are shown in Figure 2 and Figure 4, respectively.

The change from baseline to Month 9 in modified body mass index nominally favored AMVUTTRA [Table 3].

Table 3: Clinical Efficacy Results (Comparison of AMVUTTRA Treatment in Study 1 to an External Placebo Control*)

Endpoint†	Baseline, Mean (SD)		Change from Baseline to Month 9, LS Mean (SEM)		AMVUTTRA-Placebo* Treatment Difference, LS Mean (95% CI)	p-value
	AMVUTTRA N=122 (Study 1)	Placebo* N=77 (NCT01960348)	AMVUTTRA (Study 1)	Placebo* (NCT01960348)		
mNIS+7‡	60.6 (36.0)	74.6 (37.0)	-2.2 (1.4)	14.8 (2.0)	-17.0 (-21.8, -12.2)	p<0.001
Norfolk QoL-DN‡	47.1 (26.3)	55.5 (24.3)	-3.3 (1.7)	12.9 (2.2)	-16.2 (-21.7, -10.8)	p<0.001
10-meter walk test (m/sec)§	1.01 (0.39)	0.79 (0.32)	0 (0.02)	-0.13 (0.03)	0.13 (0.07, 0.19)	p<0.001
mBMI¶	1058 (234)	990 (214)	7.6 (7.9)	-60.2 (10.1)	67.8 (43.0, 92.6)	p<0.001

CI = confidence interval; LS mean = least squares mean; mBMI = modified body mass index; mNIS = modified Neuropathy Impairment Score; QoL-DN = Quality of Life-Diabetic Neuropathy; SD = standard deviation; SEM = standard error of the mean

*External placebo group from another randomized controlled trial (NCT01960348)


†All endpoints analyzed using the analysis of covariance (ANCOVA) with multiple imputation (MI) method

‡A lower number indicates less impairment/fewer symptoms

§A higher number indicates less disability/less impairment

¶mBMI: nominal p-value; body mass index (BMI; kg/m²) multiplied by serum albumin (g/L).

Figure 1: Change from Baseline in mNIS+7 (Comparison of AMVUTTRA Treatment in Study 1 to an External Placebo Control*)




A decrease in mNIS+7 indicates improvement

Δ indicates between-group treatment difference, shown as the LS mean difference (95% CI) for AMVUTTRA – placebo


*External placebo group from another randomized controlled trial (NCT01960348)

Figure 2: Histogram of mNIS+7 Change from Baseline at Month 9 (Comparison of AMVUTTRA Treatment in Study 1 to an External Placebo Control*)




Categories are mutually exclusive; patients who died before 9 months are summarized in the "Death" category only
 *External placebo group from another randomized controlled trial (NCT01960348)

Figure 3: Change from Baseline in Norfolk QoL-DN Total Score (Comparison of AMVUTTRA Treatment in Study 1 to an External Placebo Control*)



A decrease in Norfolk QoL-DN score indicates improvement
 Δ indicates between-group treatment difference, shown as the LS mean difference (95% CI) for AMVUTTRA – placebo
 *External placebo group from another randomized controlled trial (NCT01960348)

Figure 4: Histogram of Norfolk QoL-DN Total Score Change from Baseline at Month 9 (Comparison of AMVUTTRA Treatment in Study 1 to an External Placebo Control*)



Patients receiving AMVUTTRA in Study 1 experienced similar improvements relative to those in the external placebo group in mNIS+7 and Norfolk QoL-DN total score across all subgroups including age, sex, race, region, NIS score, Val30Met genotype status, and disease stage.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

AMVUTTRA is a sterile, preservative-free, clear, colorless-to-yellow solution for subcutaneous injection. AMVUTTRA is supplied as 25 mg/0.5 mL solution in a single-dose 1-mL prefilled syringe made from Type I glass with stainless steel 29-gauge needle with a needle shield. The prefilled syringe components are not made with natural rubber latex.

AMVUTTRA is available in cartons containing one single-dose prefilled syringe each.

The NDC is: 71336-1003-1.

16.2 Storage and Handling

Store at 2°C to 30°C (36°F to 86°F) in the original carton, until ready for use. Do not freeze.

17 PATIENT COUNSELING INFORMATION

Recommended Vitamin A Supplementation

Inform patients that AMVUTTRA treatment leads to a decrease in serum vitamin A levels. Instruct patients to take the recommended daily allowance of vitamin A. Advise patients to contact their healthcare provider if they experience ocular symptoms suggestive of vitamin A deficiency (e.g., night blindness) and refer them to an ophthalmologist if they develop these symptoms [see [Warnings and Precautions \(5.1\)](#)].

Pregnancy

Instruct patients that if they are pregnant or plan to become pregnant while taking AMVUTTRA they should inform their healthcare provider. Inform patients of the potential risk to the fetus, including that AMVUTTRA treatment leads to a decrease in serum vitamin A levels [see [Use in Specific Populations \(8.1\)](#) and [Clinical Pharmacology \(12.2\)](#)].

Manufactured for: Alnylam Pharmaceuticals, Inc., Cambridge, MA 02142

AMVUTTRA is a pending trademark of Alnylam Pharmaceuticals, Inc.